

VU Research Portal

Spontaneous Rayleigh-Brillouin Scattering in Molecular Gases

Wang, Y.

2019

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Wang, Y. (2019). *Spontaneous Rayleigh-Brillouin Scattering in Molecular Gases*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Contents

1	Introduction	1
1.1	Prologue	1
1.2	Theory of Rayleigh-Brillouin scattering	2
1.3	Rayleigh-Brillouin lineshape	6
1.4	Models for the Rayleigh-Brillouin lineshape	10
1.4.1	The dilute regime	12
1.4.2	The continuum regime	17
1.5	Outline of this thesis	18
2	Experimental setup and procedures	21
2.1	Rayleigh-Brillouin scattering setup with 403 nm	22
2.2	Rayleigh-Brillouin scattering setup with 532 nm	22
2.2.1	Scattering unit	24
2.2.2	Detection unit	26
2.3	Experimental data processing	33
3	Rayleigh-Brillouin scattering in SF₆ in the kinetic regime	39
3.1	Introduction	40
3.2	Experimental	43
3.3	Results	44
3.3.1	Comparison with the Tenti-S6 model	45
3.3.2	Comparison with the rough-sphere model	47
3.3.3	Comparison with the hydrodynamic model	48
3.3.4	Difference between experiment and models	48
3.4	Conclusion	49
4	Rayleigh-Brillouin light scattering spectroscopy of N₂O	51
4.1	Introduction	52
4.2	Experiment	53
4.3	Results	55
4.3.1	The Tenti-S6 model	57

4.3.2	The Grad's six-moment kinetic model	58
4.3.3	The Hammond-Wiggins hydrodynamic model	60
4.3.4	The 'rough-sphere' model	61
4.4	Discussion and Conclusion	62
5	Bulk viscosity of CO₂ from RB-spectroscopy at 532 nm	65
5.1	Introduction	66
5.2	Experiment	68
5.3	RB-scattering and line shape models	70
5.3.1	The Tenti model	71
5.3.2	The Hammond-Wiggins model	72
5.4	Results and Discussion	72
5.4.1	Measurements: Light scattering in CO ₂	72
5.4.2	Comparison with the two models	74
5.4.3	Bulk viscosities	75
5.5	Discussion and conclusion	77
6	Rayleigh-Brillouin scattering of binary mixture gases	81
6.1	Introduction	82
6.2	Model description	84
6.3	Experimental setup and results	86
6.3.1	SF ₆ and He	88
6.3.2	SF ₆ and D ₂	89
6.3.3	SF ₆ and H ₂	90
6.3.4	CO ₂ and He	92
6.3.5	CO ₂ and D ₂	93
6.3.6	Mixtures with CH ₄	94
6.4	Conclusion	95
A	Matlab code for Tenti-S6 model	99
B	Matlab code for rough-sphere model	105
C	Matlab code for Grad's six-moment model	109
D	Matlab code for HW-hydrodynamic model	113
	Bibliography	117
	Summary	129
	Samenvatting	131

CONTENTS

summary (Chinese)	135
Acknowledgements	137