

VU Research Portal

Cardiovascular events in type 2 diabetes

Becker, A.; Bos, G.; de Vegt, F.; Kostense, P.J.; Dekker, J.M.; Nijpels, G.; Heine, R.J.; Bouter, L.M.; Stehouwer, C.D.A.

published in

European Heart Journal
2003

DOI (link to publisher)

[10.1016/S0195-668X\(03\)00197-0](https://doi.org/10.1016/S0195-668X(03)00197-0)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Becker, A., Bos, G., de Vegt, F., Kostense, P. J., Dekker, J. M., Nijpels, G., Heine, R. J., Bouter, L. M., & Stehouwer, C. D. A. (2003). Cardiovascular events in type 2 diabetes: comparison with nondiabetic individuals without and with prior cardiovascular disease. 10-year follow-up of the Hoorn Study. *European Heart Journal*, 24(15), 1406-1413. [https://doi.org/10.1016/S0195-668X\(03\)00197-0](https://doi.org/10.1016/S0195-668X(03)00197-0)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

VARIANCE AND DISSENT

Reply

Clinimetrics and psychometrics: two sides of the same coin

Henrica C.W. de Vet*, Caroline B. Terwee, Lex M. Bouter

Institute for Research in Extramural Medicine, VU University Medical Center, Van der Boechorststraat 7 1081 BT, Amsterdam, The Netherlands

Accepted 15 August 2003

In this issue of the *Journal of Clinical Epidemiology*, Streiner advocates the abolishment of clinimetrics [1]. We have a number of arguments to keep clinimetrics alive and even to promote its use. The merit of Feinstein's introduction of the term "Clinimetrics" and the appearance of his book on "Clinimetrics" [2] is the increased attention for measurement issues within medicine. We believe that this would never have occurred with the introduction of psychometry in medicine, which clinicians would easily disqualify as being none of their business. Apart from the need to use terms attractive to clinicians, the approaches used should fit in the frame of thinking of doctors; for example, to assess interobserver variation clinicians prefer the Bland and Altman method [3], which expresses the measurement error in the dimension of measurement, while psychologists would prefer an intraclass correlation coefficient (ICC), a number between 0 and 1 [4]. Such a coefficient is less attractive to clinicians because it does not give them a clue about the size of the measurement error. By the same token, indexes based on clinical judgement fit more in the mode of thinking of clinicians than the construction of scales using sophisticated statistical methods [2].

There obviously exists a substantial overlap between clinimetrics, psychometrics, and biometrics. Were these terms to be defined anew, we would propose the term "metrics" as an indication of a measurement discipline and label it as biometrics, psychometrics, or clinimetrics, depending on the field of application. Feinstein [2] and Fayers and Hand [5] propose to reserve psychometrics for unidimensional scales and clinimetrics for multidimensional indexes. We agree with Fayers and Hand that the clinimetric and the psychometric approach are not contradictory, but serve different aims: the clinimetric approach is directed at the development of instruments to measure multiple constructs with a single index (e.g., the Apgar score or the TNM score to assess tumors, nodes and metastases in cancer patients), while the psychometric approach is appropriate to develop

instruments that measure a single construct using multiple items (e.g., depression or anxiety) [5]. Fayers and Hand elegantly explain this appropriateness for different purposes by distinguishing two different types of variables to be included in measurement instruments: indicator variables and causal variables. Indicator variables are variables that correlate with the underlying construct to be measured, but do not alter or influence this construct. Causal variables influence the construct to be measured by a causal relationship, for example, being symptoms of the disease or side effects of the treatment. In general, psychometric measurement instruments only include indicator variables that are always correlated with the construct under study, while clinimetric instruments may also include causal variables. The difference between the clinimetric and the psychometric approach is most evident in the development phase of the instrument. In psychology, unidimensional constructs (such as intelligence, personality, or internal locus of control) are often assessed, and for this purpose psychometric methods are needed. Factor analysis, to examine the underlying dimensions of the scale, and assessments of the internal consistency of the scale or subscales are important psychometric properties. In medicine, composite indexes combining different symptoms and characteristics are more often at issue, and for this purpose clinimetric methods are appropriate. These clinimetric and psychometric approaches typically meet in the construction of measurements to measure quality of life, for which both clinical and psychological phenomena are important, and both indicator and causal variables may be included in parts of the instrument. In the evaluation of measurement instruments, when assessing validity, reproducibility, or responsiveness, the differences between clinimetric and psychometric approaches are less obvious and the characteristics can be considered as clinimetric or psychometric, depending on the measurement of clinical or psychologic phenomenon.

With Feinstein [2], others see differences between clinimetrics and psychometrics in the way the instruments are developed (more content driven in case of clinimetrics, more statistically driven in case of psychometrics) [5–7]. Comparison of both approaches showed that different items would be

* Corresponding author. Tel. +31 20 444 8176; fax: +31 20 444 8181.
E-mail address: HCV.deVet@vumc.nl (H.C.W. de Vet).

included in an instrument depending on the method used [6,7]. This indicates that the approaches are different.

As Streiner [1] correctly points out, none of these distinctions, either based on fields of application, number of dimensions, or the principle of item selection are exclusive. It is a matter of degrees: for clinical purposes, the instruments are more often multidimensional and typically constructed on the basis what patients or clinicians consider to be important. This indeed hampers a clear distinction between the terms psychometrics and clinimetrics. Both metric disciplines make use of the same methodologic and statistical approaches, depending on the goal and the subject of measurement. The remark of Streiner [1] that clinimetrists are ignoring the literature on psychometrics can easily be refuted by various examples. Disability, a clinical phenomenon, has been examined by techniques from the item response theory [8,9], and the generalizability theory has been applied to range of motion measurements [10].

Most clinimetrists, including Feinstein, are very much aware of the merits of psychometrics for the development of clinimetrics. In fact, the book of Streiner and Norman on Health Measurement Scales, which is written from a psychometric perspective, is widely used in clinical and health research as well as education. It has greatly helped to prevent clinimetrists from being ignorant.

We strongly believe that clinimetrics is here to stay. The term is instrumental in involving medical doctors in metrics. Furthermore, its vocabulary rightly stresses the aspects of

metrics, which are most important for measurement in clinical research and practice.

References

- [1] Streiner D. Clinimetrics vs. psychometrics: an unnecessary distinction. *J Clin Epidemiol* 2003;56:1142–45.
- [2] Feinstein AR. *Clinimetrics*. New Haven, CT: Yale University Press; 1987.
- [3] Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurements. *Lancet* 1986;i: 307–10.
- [4] Streiner DL, Norman GR. *Health measurement scales. A practical guide to their development and use*. Oxford: Oxford University Press; 1995.
- [5] Fayers PM, Hand DJ. Causal variables, indicator variables and measurement scales: an example from quality of life. *J R Stat Soc* 2002; 165:233–61.
- [6] Juniper EF, Guyatt G, Streiner DL, King DR. Clinical impact versus factor analysis for quality of life questionnaire construction. *J Clin Epidemiol* 1997;50:233–8.
- [7] Marx RG, Bombardier C, Hogg-Johnson S, Wright JG. Clinimetric and psychometric strategies for development of a health measurement scale. *J Clin Epidemiol* 1999;52:105–11.
- [8] Garrett AM. RASCH analysis of the Roland Disability Questionnaire. *Spine* 2003;28:79–84.
- [9] White LJ, Velozo CA. The use of RASCH measurement to improve the Oswestry classification scheme. *Arch Phys Med Rehabil* 2002; 83:822–31.
- [10] Roebroeck MW, Harlaar J, Lankhorst GJ. Reliability assessment of isometric knee extension measurements with a computer-assisted hand-held dynamometer. *Arch Phys Med Rehabil* 1998;79:442–8.